

# Candidate Marks Report

*Series : 6 2018*

This candidate's script has been assessed using On-Screen Marking. The marks are therefore not shown on the script itself, but are summarised in the table below.

---

Centre No :	Assessment Code :	H432
Candidate No :	Component Code :	01
Candidate Name :		

---

Total Marks :

In the table below 'Total Mark' records the mark scored by this candidate.  
'Max Mark' records the Maximum Mark available for the question.

## SECTION A

You should spend a maximum of 20 minutes on this section.

Write your answer to each question in the box provided.

Answer all the questions.

- 1 A sample of boron contains the isotopes  $^{10}\text{B}$  and  $^{11}\text{B}$ . 5. elec  
The relative atomic mass of the boron sample is 10.8.

What is the percentage of  $^{11}\text{B}$  atoms in the sample of boron?

- A 8.0%  
B 20%  
C 80%  
D 92%

$$10.8 = \frac{(10 \times 30) + (11 \times 80)}{100}$$

Your answer

[1]

- 2 In the compound  $[\text{ICl}_2]^+ [\text{SbCl}_6]^-$ , the oxidation number of chlorine is  $-1$ .

What are the oxidation numbers of I and Sb in the compound?

	I	Sb
A	+1	+5
B	+1	+7
C	+3	+5
D	+3	+7

Your answer

[1]

- 3 What is the number of hydrogen atoms in 0.125 mol of  $\text{C}_2\text{H}_5\text{OH}$ ?


- A  $7.525 \times 10^{22}$  X  
B  $4.515 \times 10^{23}$  ✓  
C  $3.7625 \times 10^{23}$   
D  $3.612 \times 10^{24}$

$\frac{6}{9}$

Your answer

[1]





3

4 A student titrates a standard solution of barium hydroxide,  $\text{Ba(OH)}_2$ , with nitric acid,  $\text{HNO}_3$ .  
 25.00  $\text{cm}^3$  of  $0.0450 \text{ mol dm}^{-3}$   $\text{Ba(OH)}_2$  are needed to neutralise 23.35  $\text{cm}^3$  of  $\text{HNO}_3(\text{aq})$ .

What is the concentration, in  $\text{mol dm}^{-3}$ , of the nitric acid?

- A 0.0241
- B 0.0482
- C 0.0900
- D 0.0964

$$1.125 \times 10^{-3} \text{ Ba(OH)}_2$$


$$\underline{2.25 \times 10^{-3} \text{ mol}}$$

$$0.0964$$

Your answer

**D**

[1]



5 Which statement best explains why nitrogen has a larger first ionisation energy than oxygen?

- A N atoms have less repulsion between p-orbital electrons than O atoms. ✓
- B N atoms have a smaller nuclear charge than O atoms. ✗
- C N atoms lose an electron from the 2s subshell, while O atoms lose an electron from the 2p subshell. ✗
- D N atoms have an odd number of electrons, while O atoms have an even number.

Your answer


**A**

[1]

6 In the Periodic Table, element X is in Group 2 and element Y is in Group 15 (5).

What is the likely formula of an ionic compound of X and Y?

- A  $\text{X}_2\text{Y}_5$
- B  $\text{X}_2\text{Y}_3$
- C  $\text{X}_3\text{Y}_2$
- D  $\text{X}_5\text{Y}_2$




Your answer

**C**

[1]





7 Which statement about ammonium carbonate is **not** correct?

- A It reacts with  $\text{Ba}(\text{NO}_3)_2(\text{aq})$  to form a white precipitate. ~~✓~~
- B It effervesces with dilute nitric acid. ✗
- C It releases an alkaline gas with warm  $\text{NaOH}(\text{aq})$ . ✗
- D It has the formula  $\text{NH}_4\text{CO}_3$ . ✓





Your answer

D

[1]

8 A reaction is first order with respect to a reactant X.

Which rate-concentration graph for reactant X is the correct shape?




Your answer

A

[1]



- 9 The reversible reaction of sulfur dioxide and oxygen to form sulfur trioxide is shown below.



An equilibrium mixture contains 2.4 mol  $\text{SO}_2$ , 1.2 mol  $\text{O}_2$  and 0.4 mol  $\text{SO}_3$ .  
The total pressure is 250 atm.

What is the partial pressure of  $\text{SO}_3$ ?

$$\frac{0.4}{4} = 0.1$$

$$\times 250$$

- A 15 atm  
B 25 atm  
C 100 atm  
D 200 atm

Your answer

B

[1]

- 10 A buffer solution is prepared by mixing 200 cm<sup>3</sup> of 2.00 mol dm<sup>-3</sup> propanoic acid,  $\text{CH}_3\text{CH}_2\text{COOH}$ , with 600 cm<sup>3</sup> of 1.00 mol dm<sup>-3</sup> sodium propanoate,  $\text{CH}_3\text{CH}_2\text{COONa}$ .

$K_a$  for  $\text{CH}_3\text{CH}_2\text{COOH} = 1.32 \times 10^{-5} \text{ mol dm}^{-3}$

What is the pH of the buffer solution?

- A 4.58  
B 4.70  
C 5.06  
D 5.18

$$\frac{200}{1000} \times 2 = 0.4 \text{ mol HA}$$

$$0.6 \text{ mol A}^-$$

$$\frac{0.4}{0.8} = 0.5 \text{ HA}$$

$$\frac{0.6}{0.8} = 0.75 \text{ A}^-$$

Your answer

C

[1]

$$\Rightarrow K_a \times \frac{[\text{HA}]}{[\text{A}^-]} = [\text{H}^+]$$


$$-\log [\text{H}^+]$$



- 11 The table below shows standard entropies,  $S^\ominus$ .

Substance	CO(g)	H <sub>2</sub> (g)	CH <sub>3</sub> OH(l)
$S^\ominus/\text{J mol}^{-1}\text{K}^{-1}$	197.6	130.6	239.7

What is the entropy change,  $\Delta S^\ominus$ , in  $\text{J mol}^{-1}\text{K}^{-1}$ , for the following reaction?



- A -219.1  
B -88.5  
C +88.5  
D +219.1

$$239.7 - (197.6 + 2(130.6))$$

Your answer

A

[1]

- 12 The redox equilibria for a hydrogen-oxygen fuel cell in alkaline solution are shown below.




What is the equation for the overall cell reaction?


- A  $\text{H}_2\text{(g)} + 4\text{OH}^-\text{(aq)} \rightarrow 3\text{H}_2\text{O(l)} + \frac{1}{2}\text{O}_2\text{(g)}$   
B  $3\text{H}_2\text{O(l)} + \frac{1}{2}\text{O}_2 \rightarrow \text{H}_2\text{(g)} + 4\text{OH}^-\text{(aq)}$   
C  $\text{H}_2\text{O(l)} \rightarrow \text{H}_2\text{(g)} + \frac{1}{2}\text{O}_2\text{(g)}$   
D  $\text{H}_2\text{(g)} + \frac{1}{2}\text{O}_2\text{(g)} \rightarrow \text{H}_2\text{O(l)}$

Your answer

D



[1]



13 Which enthalpy change(s) is/are endothermic?

break bond

- 1 The bond enthalpy of the C-H bond
- 2 The second electron affinity of oxygen
- 3 The standard enthalpy change of formation of magnesium

- A 1, 2 and 3
- B Only 1 and 2
- C Only 2 and 3
- D Only 1

By down

Your answer:

C

[1]

14 Which statement(s) explain(s) why reaction rates increase as temperature increases?

- 1 The activation energy is less.
- 2 Collisions between molecules are more frequent.
- 3 A greater proportion of molecules have energy greater than the activation energy.

- A 1, 2 and 3
- B Only 1 and 2
- C Only 2 and 3
- D Only 1

Your answer:

C

[1]



15 Which statement(s) is/are correct for the complex  $\text{Pt}(\text{NH}_3)_2\text{Cl}_2$ ?

- 1 One of its stereoisomers is used as an anti-cancer drug. ✓  
2 It has bond angles of  $109.5^\circ$ . ✗  
3 It has optical isomers. ✗

- A 1, 2 and 3  
B Only 1 and 2  
C Only 2 and 3  
D Only 1

Your answer

D

[1]





BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE



Answer all the questions.

16 This question is about enthalpy changes.

(a) Table 16.1 shows enthalpy changes that can be used to determine the enthalpy change of hydration of fluoride ions,  $F^-$ .

Enthalpy change	Energy / $\text{kJ mol}^{-1}$
Hydration of $\text{Ca}^{2+}$	-1609
Solution of $\text{CaF}_2$	+13
Lattice enthalpy of $\text{CaF}_2$	-2630

Table 16.1


(i) Explain what is meant by the term *enthalpy change of hydration*.

the enthalpy change when gaseous aqueous ions  
are formed from 1 mol of water reacts with  
gaseous ions to form aqueous ions

[2]

(ii) The enthalpy change of hydration of  $F^-$  can be determined using the enthalpy changes in Table 16.1 and the incomplete energy cycle below.

On the dotted lines, add the species present, including state symbols.



[4]



- (iii) Calculate the enthalpy change of hydration of fluoride ions,  $F^-$ .

$$a + d = b + c$$

$$a - 1609 = -2630 + 13$$

$$a = -2630 + 1609 + 13$$

$$-1008 \div 2 = -504$$

$$-504$$

enthalpy change of hydration = ~~1008~~  $-504$   $\text{kJ mol}^{-1}$  [2]

- (iv) Predict how the enthalpy changes of hydration of  $F^-$  and  $Cl^-$  would differ.

Explain your answer.

enthalpy change of  $Cl^-$  would be lower as  
 $Cl$  ~~be~~ bond weaker, ~~breaks down~~ breaks and  
 dissolves easier.


[2]

$$-1516 = \frac{1008}{2}$$

$$+1609$$



(b) Fluorine reacts with steam as shown in the equation below.



Average bond enthalpies are shown in the table.

Bond	Average bond enthalpy / $\text{kJ mol}^{-1}$
O-H	+464
O=O	+498
H-F	+568

(i) Explain what is meant by the term *average bond enthalpy*.

relative amount of energy required to break a bond

[2]

(ii) Calculate the bond enthalpy of the F-F bond.

$$\begin{aligned} & (2\text{F-F} + 4\text{O-H}) - (\text{O=O} + 4\text{H-F}) \\ & = (2\text{F-F} + 1856) - (2770) \end{aligned}$$

$$2\text{F-F} - 914 = -598$$

$$2\text{F-F} = 316$$

$$\text{F-F} = 158$$

bond enthalpy = 158  $\text{kJ mol}^{-1}$  [3]



BLANK PAGE


2172 - 2770

PLEASE DO NOT WRITE ON THIS PAGE



17 This question is about reaction rates.

Aqueous iron(III) ions,  $\text{Fe}^{3+}(\text{aq})$ , react with aqueous iodide ions,  $\text{I}^{-}(\text{aq})$  as shown below.



A student carries out three experiments to investigate how different concentrations of  $\text{Fe}^{3+}(\text{aq})$  and  $\text{I}^{-}(\text{aq})$  affect the initial rate of this reaction. The results are shown below.

Experiment	$[\text{Fe}^{3+}(\text{aq})]$ / $\text{mol dm}^{-3}$	$[\text{I}^{-}(\text{aq})]$ / $\text{mol dm}^{-3}$	Initial rate / $\text{mol dm}^{-3} \text{s}^{-1}$
1	$4.00 \times 10^{-2}$	$3.00 \times 10^{-2}$	$8.10 \times 10^{-4}$
2	$8.00 \times 10^{-2}$	$3.00 \times 10^{-2}$	$1.62 \times 10^{-3}$
3	$4.00 \times 10^{-2}$	$6.00 \times 10^{-2}$	$3.24 \times 10^{-3}$

(a)\* Determine the rate constant and a possible two-step mechanism for this reaction that are consistent with these results. [6]




$\text{Fe}^{3+}$  is first order and  $\text{I}^{-}$  is second order

$$\text{rate} = k [\text{Fe}^{3+}] [\text{I}^{-}]^2 \quad \frac{\text{rate}}{[\text{Fe}^{3+}] [\text{I}^{-}]^2} = k$$

$$\frac{8.1 \times 10^{-4}}{(4 \times 10^{-2})(3 \times 10^{-2})^2} = k = 22.5$$

$$\frac{\text{mol dm}^{-3} \text{s}^{-1}}{(\text{mol dm}^{-3})(\text{mol dm}^{-3})(\text{mol dm}^{-3})}$$

$$\text{rate constant} = \text{dm}^6 \text{mol}^{-2} \text{s}^{-1} = 22.5 \text{ dm}^6 \text{mol}^{-2} \text{s}^{-1}$$



Additional answer space if required



.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....



- (b) A student carries out an investigation to find the activation energy,  $E_a$ , and the pre-exponential factor,  $A$ , of a reaction.

The student determines the rate constant,  $k$ , at different temperatures,  $T$ . The student then plots a graph of  $\ln k$  against  $1/T$  as shown below.





$$\ln A + \frac{E_a}{R} \times \frac{1}{T}$$

+c  
gradient 2c

- (i) Draw a best-fit straight line and calculate the activation energy, in  $\text{J mol}^{-1}$ .  
Give your answer to **three** significant figures.

8.314

Show your working.

$$\frac{-4}{5 \times 10^{-3}} = -800$$

$$-800 = -\frac{E_a}{R}$$

$$-800 \times -8.314 = E_a$$

activation energy,  $E_a = +$  ..... 6.650 .....  $\text{J mol}^{-1}$  [3]

- (ii) Use the graph to calculate the value of the pre-exponential factor, A.


Show your working.

$$\ln k = 31.3$$

$$k = e^{31.3}$$

pre-exponential factor, A = ..... ~~3.92 x 10<sup>12</sup>~~ ..... [2]  
3.92 x 10<sup>13</sup>

18 Nitrogen monoxide,  $\text{NO}$ , and oxygen,  $\text{O}_2$ , react to form nitrogen dioxide,  $\text{NO}_2$ , in the reversible reaction shown in **equilibrium 18.1**.




(a) Write an expression for  $K_c$  for this equilibrium and state the units.

$$K_c = \frac{[\text{NO}_2]^2}{[\text{NO}]^2 [\text{O}_2]}$$

$$\text{Units} = \dots \text{dm}^3 \text{mol}^{-1} \dots$$

[2]

(b) A chemist mixes together nitrogen and oxygen and pressurises the gases so that their total gas volume is  $4.0 \text{ dm}^3$ .



- The mixture is allowed to reach equilibrium at constant temperature and volume.
- The equilibrium mixture contains  $0.40 \text{ mol NO}$  and  $0.80 \text{ mol O}_2$ .
- Under these conditions, the numerical value of  $K_c$  is  $45$ .

Calculate the amount, in mol, of  $\text{NO}_2$  in the equilibrium mixture.

	$\text{N}_2$	$\frac{3}{2}\text{O}_2$	$\rightleftharpoons$	$2\text{NO}$	$2\text{NO}_2$
initial mol		1.4		0	0
equilibrium mol		0.8		0.4	0.4
change	-0.4	-0.4		+0.4	+0.4
balance	-0.4	-0.6		+0.4	+0.4



$$0.3 \times 4$$


initial			
equilibrium	0.4	0.8	1.2
change			
balance			
volume	4	4	4
conc	0.1	0.2	0.3

amount of  $\text{NO}_2 = \dots\dots\dots 1.2 \dots\dots\dots \text{mol}$  [4]

$$45 = \frac{[\text{NO}_2]^2}{(0.1)^2 (0.2)} \quad \dots \quad 0.3 = [\text{NO}_2]$$



(c) The values of  $K_p$  for equilibrium 18.1 at 298K and 1000K are shown below.



Equilibrium 18.1

Temperature / K	$K_p / \text{atm}^{-1}$
298	$K_p = 2.19 \times 10^{12}$
1000	$K_p = 2.03 \times 10^{-1}$

increase in temp  
shifts to endothermic side  
•  $K_c$  decreases, e-5

(i) Predict, with a reason, whether the forward reaction is exothermic or endothermic.


..... forward reaction is exothermic as ~~the~~ when temp. increases equilibrium shifts to endothermic side which is ..... [1]  
towards reactants as value of  $K_p$  decreases

(ii) The chemist increases the pressure of the equilibrium mixture at the same temperature.

State, and explain in terms of  $K_p$ , how you would expect the equilibrium position to change.

..... value of  $K_p$  stays the same. Increase in pressure shifts equilibrium to the right towards the products as that's the side with fewer moles. ....  
or when you increase pressure  $K_p$  <sup>ratio</sup> (reactants) ~~ratio~~ denominator increases so ~~to~~ to oppose that change and keep value of  $K_p$  the same, increase pressure of numerator (products) and decrease denominator. .... [3]  
(reactants)





19 This question is about acids and bases found in the home.

(a) Ethanoic acid, CH<sub>3</sub>COOH, is the acid present in vinegar. *vinegar contains acid*

A student carries out an experiment to determine the pK<sub>a</sub> value of CH<sub>3</sub>COOH.

- The concentration of CH<sub>3</sub>COOH in the vinegar is 0.870 mol dm<sup>-3</sup>
- The pH of the vinegar is 2.41

(i) Write the expression for the acid dissociation constant, K<sub>a</sub>, of CH<sub>3</sub>COOH.

$$K_a = \frac{[\text{CH}_3\text{COO}^-][\text{H}^+]}{\text{CH}_3\text{COOH}}$$

[1]

(ii) Calculate the pK<sub>a</sub> value of CH<sub>3</sub>COOH.

Give your answer to **two** decimal places.

$$10^{-2.41} = [\text{H}^+] = 3.89 \times 10^{-3}$$

$$K_a = \frac{[\text{H}^+]^2}{\text{HA}} = \frac{(3.89 \times 10^{-3})^2}{0.87} = 1.74 \times 10^{-5}$$

$$-\log(1.74 \times 10^{-5}) = \text{pH}$$

pK<sub>a</sub> = 4.76 [3]

(iii) Determine the percentage dissociation of ethanoic acid in the vinegar.

Give your answer to **three** significant figures.

$$\frac{3.89 \times 10^{-3}}{0.87} \times 100$$

percentage dissociation = 0.447 % [1]



(b) Many solid drain cleaners are based on sodium hydroxide,  $\text{NaOH}$ .

- A student dissolves  $1.26\text{ g}$  of a drain cleaner in water and makes up the solution to  $100.0\text{ cm}^3$ .
- The student measures the pH of this solution as  $13.48$ .  $K_w = 1 \times 10^{-14}$

Determine the percentage, by mass, of  $\text{NaOH}$  in the drain cleaner.

Give your answer to three significant figures.

$$K_w = [\text{OH}^-][\text{H}^+] \quad 10^{-13.48} = [\text{H}^+] = 3.31 \times 10^{-14}$$

$$\frac{1 \times 10^{-14}}{3.31 \times 10^{-14}} = [\text{OH}^-] = 0.302 \text{ mol dm}^{-3}$$

$$0.302 \times \frac{100}{1000} = 0.0302 \text{ mol NaOH}$$



$$0.0302 \times 40 = 1.208 \text{ g}$$

$$\frac{1.208}{1.26} \times 100 = 95.9\%$$

percentage = 95.9 % [4]


(c) Sodium carbonate,  $\text{Na}_2\text{CO}_3$  is a base used in washing soda.

$\text{Na}_2\text{CO}_3$  contains the carbonate ion,  $\text{CO}_3^{2-}$ , shown below.



Draw the 'dot-and-cross' diagram for the carbonate ion.

Show outer electrons only and use different symbols for electrons from C and O and any 'extra' electrons.



$\rightarrow = \text{extra electron}$

[2]

Turn over



20 This question is about the halogen group of elements and some of their compounds.

(a) The halogens show trends in their properties down the group.

The boiling points of three halogens are shown below.


Halogen	Boiling point/°C
Chlorine	-35
Bromine	59
Iodine	184

Explain why the halogens show this trend in boiling points.

As you go down the halogen group, boiling point increases as the number of electrons increases and stronger London forces. Stronger London forces requires the use of more energy to overcome / break.

[3]


(b) Hydrogen iodide, HI, is decomposed by heat into its elements:



The decomposition is much faster in the presence of a platinum catalyst.

Complete the enthalpy profile diagram for this reaction using formulae for the reactants and products.


- Use  $E_a$  to label the activation energy without a catalyst.
- Use  $E_c$  to label the activation energy with a catalyst.
- Use  $\Delta H$  to label the enthalpy change of reaction.



[3]



simple



1 mol = 24000 cm<sup>3</sup>

(c) Compound A is an oxide of chlorine that is a liquid at room temperature and pressure and has a boiling point of 83°C.

When 0.4485 g of A is heated to 100°C at  $1.00 \times 10^5$  Pa, 76.0 cm<sup>3</sup> of gas is produced.

Determine the molecular formula of compound A.

Show all your working.

$PV = nRT$

$\frac{PV}{RT} = n$


$(1 \times 10^5) \times (7.6 \times 10^{-5})$

$8.314 \times 373$

$= 2.45 \times 10^{-3} \text{ mol}$

$\frac{0.4485}{2.45 \times 10^{-3}} = 183 = nR$

$\frac{183}{87} = 2.1$



molecular formula of A =  $\text{Cl}_2\text{O}_7$  [4]



- (d) Compound B is an iodate(V) salt of a Group 1 metal.  
The iodate(V) ion has the formula  $\text{IO}_3^-$ .




A student carries out a titration to find the formula of compound B.

**Step 1:** The student dissolves 1.55 g of B in water and makes up the solution to 250.0 cm<sup>3</sup> in a volumetric flask.


$\times 10$

**Step 2:** The student pipettes 25.00 cm<sup>3</sup> of the solution of B into a conical flask, followed by 10 cm<sup>3</sup> of dilute sulfuric acid and an excess of KI(aq).

The iodate(V) ions are reduced to iodine, as shown below.



**Step 3:** The resulting mixture is titrated with 0.150 mol dm<sup>-3</sup>  $\text{Na}_2\text{S}_2\text{O}_3(\text{aq})$ .



The student repeats **step 2** and **step 3** until concordant titres are obtained.

**Titration readings**

Titration	Trial	1	2	3
Final burette reading / cm <sup>3</sup>	24.00	47.40	23.75	47.05
Initial burette reading / cm <sup>3</sup>	0.00	24.00	0.00	23.20
Titre / cm <sup>3</sup>	24.00	23.40	23.75	23.85

Table 20.1

- (i) Complete **Table 20.1** and calculate the mean titre that the student should use for analysing the results.

$$\frac{23.85 + 23.75}{2} = 23.80$$

mean titre = ..... 23.80 ..... cm<sup>3</sup> [2]

- (ii) The uncertainty in each burette reading is  $\pm 0.05 \text{ cm}^3$ .

Calculate the percentage uncertainty in the titre obtained from titration 1.

Give your answer to two decimal places.

$$\frac{0.05 \times 2}{23.40} \times 100$$

percentage uncertainty = ..... 0.43 ..... % [1]





- (iii) Describe and explain how the student should determine the end point of this titration accurately.

... I<sub>2</sub> in a beaker... Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> in a burette... Open the tap of the burette and allow Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> to <sup>pour</sup> ~~run~~ into I<sub>2</sub> &... When end point is approaching, ~~let~~ <sup>is being added</sup> ~~it~~ <sup>is</sup> adjust the tap till ~~it~~ <sup>Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution</sup> [2]  
 (iv) Determine the relative formula mass and formula of the Group 1 iodate(V), B.

Show your working.

$$\frac{23.8}{1000} \times 0.15 = 3.57 \times 10^{-3} \text{ mol Na}_2\text{S}_2\text{O}_3$$

$$\div 2 = 1.785 \times 10^{-3} \text{ mol I}_2$$

$$\div 3$$

$$= 5.95 \times 10^{-4} \text{ mol IO}_3^-$$

$$\times 10$$

$$= 5.95 \times 10^{-3} \text{ mol IO}_3^-$$

$$\frac{1.55}{5.95 \times 10^{-3}} = mR = 260.5$$

$$260.5 - 174.9 = 85.6 \approx Rb$$

relative formula mass of B = 260.5.....

formula of B = RbIO<sub>3</sub>..... [5]



21 This question is about some reactions of d block elements and their ions.

Table 21.1 shows standard electrode potentials which will be needed within this question.

$Zn^{2+}(aq) + 2e^-$	$\rightleftharpoons$	$Zn(s)$	$E^\ominus = -0.76V$	reducing
$Cr^{3+}(aq) + e^-$	$\rightleftharpoons$	$Cr^{2+}(aq)$	$E^\ominus = -0.42V$	
$Ni^{2+}(aq) + 2e^-$	$\rightleftharpoons$	$Ni(s)$	$E^\ominus = -0.25V$	reducing
$I_2(aq) + 2e^-$	$\rightleftharpoons$	$2I^-(aq)$	$E^\ominus = +0.54V$	oxidising
$Fe^{3+}(aq) + e^-$	$\rightleftharpoons$	$Fe^{2+}(aq)$	$E^\ominus = +0.77V$	
$Cr_2O_7^{2-}(aq) + 14H^+(aq) + 6e^-$	$\rightleftharpoons$	$2Cr^{3+}(aq) + 7H_2O(l)$	$E^\ominus = +1.33V$	oxidising
$H_2O_2(aq) + 2H^+(aq) + 2e^-$	$\rightleftharpoons$	$2H_2O(l)$	$E^\ominus = +1.78V$	


Table 21.1

(a) Complete the electron configuration of

a Ni atom:  $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^8$


a Ni<sup>2+</sup> ion:  $1s^2 2s^2 2p^6 3s^2 3p^6 3d^8$  [2]

(b) A standard cell is set up in the laboratory with the cell reaction shown below.



(i) Draw a labelled diagram to show how this cell could be set up to measure its standard cell potential.

Include details of apparatus, solutions and the standard conditions required.



Standard conditions  $298K, 100kPa$

[4]




(ii) Predict the standard cell potential of this cell.

standard cell potential = 0.79 V [1]

(c) Use the information in Table 21.1 to help you answer both parts of this question.

(i) Write the overall equation for the oxidation of Fe<sup>2+</sup> by acidified H<sub>2</sub>O<sub>2</sub>.




(ii) Zinc reacts with acidified Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> ions to form Cr<sup>3+</sup> ions in two stages.

Explain why this happens in terms of electrode potentials and equilibria.

Include overall equations for the reactions which occur.

Standard electrode potential for zinc is  $Zn$  has a more negative than Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> so equilibrium position of


is shifted more to the left so Zn is more likely to act as a reducing agent - reduces Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> as itself is oxidised








cell potential = 2.09

when excess of Zn added it's reduced further from Cr<sup>3+</sup> to Cr<sup>2+</sup> as standard electrode potential for zinc is still

more negative so its position of equilibrium for zinc lies to the left and acts as a reducing agent.



cell potential = 0.34



(d)\* Three different reactions of copper compounds are described below.

Reaction 1: Aqueous  $\text{CuSO}_4$  reacts with excess aqueous ammonia in a ligand substitution reaction. A deep-blue solution is formed, containing an octahedral complex ion, C, which is a trans isomer.  $[\text{Cu}(\text{H}_2\text{O})_2(\text{NH}_3)_4]^{2+}$

Reaction 2:  $\text{Cu}_2\text{O}$  reacts with hot dilute sulfuric acid in a disproportionation reaction. A blue solution, D, and a brown solid, E, are formed.


Reaction 3:  $\text{CuO}$  reacts with warm dilute nitric acid in a neutralisation reaction, to form a blue solution. Unreacted copper(II) oxide is filtered off, and the solution is left overnight in an evaporating basin.

$\text{CuO} + \text{HNO}_3 \rightarrow \text{CuNO}_3 + \text{H}_2\text{O}$   
 A hydrated salt, F, crystallises, with the percentage composition by mass: Cu, 26.29%; H, 2.48%; N, 11.59%; O, 59.63%.  
 $+ \text{CuO} \rightarrow x \text{H}_2\text{O}$


Identify C-F by formulae or structures, as appropriate.

Include equations, any changes in oxidation number, and working.

[6]



$\frac{26.29}{63.5}$	$\frac{2.48}{1}$	$\frac{11.59}{14}$	$\frac{59.63}{16}$
= 0.414	2.48	0.828	3.727
0.414	0.414	0.414	0.414
1	6	2	9



Additional answer space if required.



.....

.....

.....

.....

.....

.....

.....

.....

.....

**END OF QUESTION PAPER**



**ADDITIONAL ANSWER SPACE**

If additional space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margin(s).

A large rectangular area with horizontal dotted lines for writing, intended for providing additional answer space. A solid vertical line runs down the left side of this area, creating a margin. The dotted lines extend across the width of the page, leaving a narrow margin on the right side.





Area with horizontal dotted lines for writing.

# OCR

Oxford Cambridge and RSA

### Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website ([www.ocr.org.uk](http://www.ocr.org.uk)) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

