

Monday 20 June 2022 – Morning

A Level Chemistry B (Salters)

H433/02 Scientific literacy in chemistry

Time allowed: 2 hours 15 minutes

You must have:

- a clean copy of the Advance Notice Article (inside this document)
- the Data Sheet for Chemistry B

You can use:

- · a scientific or graphical calculator
- an HB pencil



											/
Please write clearly in black ink. Do not write in the barcodes.											
Centre number						Candidate number					
First name(s)											
Last name											

INSTRUCTIONS


- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided. If you need extra space use the lined pages at the end of this booklet. The question numbers must be clearly shown.
- · Answer all the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.

INFORMATION

- The total mark for this paper is **100**.
- The marks for each question are shown in brackets [].
- Quality of extended response will be assessed in questions marked with an asterisk (*).
- This document has 20 pages.

ADVICE

Read each question carefully before you start your answer.



Answer all the questions.

1 Portland cement is a powder that 'sets' to a strong solid when water is added. The cement is usually mixed with gravel so that it sets to make concrete.

The typical ingredients of Portland cement are shown in the table.

Ingredients (solids)	% by mass
CaO	61 – 67
SiO ₂	19 – 23
Al_2O_3	2.5 – 6
Fe ₂ O ₃	0 – 6
SO ₃	1.5 – 4.5

1		
(a)	e the systematic name for Fe ₂ O ₃ .	
(b)	(i)	A hazard warning for Portland cement states that it is alkaline.
		One of the ingredients in the table reacts with water to form an alkaline solution.
		Write an equation for the reaction of this ingredient with water including any ions that are formed.
		[2]
	(ii)	When Portland cement sets, the calcium oxide absorbs carbon dioxide from the air.
		Suggest an equation for the reaction that occurs.
		[1]
(c)		e way of testing for the presence of calcium in the cement would be a flame test, though iron present would interfere.
	(i)	What is the flame colour of calcium?
		[1]

	(ii)	The flame colour is caused by bright lines in the atomic emission spectrum of calci Explain how these lines are formed.	um.
	(iii)	Calculate the frequency of a line of wavelength 6.16×10^{-7} m.	
		frequency =	Hz [2]
(d)	Fe ₂	20 ₃ and CaO are present in Portland cement.	
	(i)	In which 'blocks' of the periodic table are these elements found?	
		Ca	
		Fe	[1]
	(ii)	The electron configuration of a calcium ion is $1s^22s^22p^63s^23p^6$.	
		How does the electron configuration of the ${\bf cation}$ in ${\rm Fe_2O_3}$ differ from this?	
	(iii)	Explain why iron can form two stable cations.	[1]
			[1]
(e)	Fe ₂	₂ O ₃ is soluble in concentrated hydrochloric acid.	
	Exc	cess sodium hydroxide is added to this solution.	
	Nar	me the precipitate formed and describe its colour.	
	nan	me:	
	cold	our.	[4]

		4
(f)	(i)	A student has a solution containing $Fe^{3+}(aq)$ and reads that it will react with iodide ions to form iodine. The iodine can then be titrated with sodium thiosulfate.
		Describe a method to measure the volume of sodium thiosulfate solution required to react with the iodine from $25\mathrm{cm}^3$ of Fe ³⁺ (aq).
		[5]
	(ii)	The student finds that $25.0\mathrm{cm^3}$ of $\mathrm{Fe^{3^+}}(aq)$ requires $3.2\mathrm{cm^3}$ of $0.510\mathrm{moldm^{-3}}$ sodium thiosulfate.
		Calculate the concentration of the Fe ³⁺ (aq).
		$2Fe^{3+} + 2I^{-} \rightarrow 2Fe^{2+} + I_{2}$
		$2S_2O_3^{2-} + I_2 \rightarrow S_4O_6^{2-} + 2I^-$
		Give your answer to an appropriate number of significant figures.

 $[Fe^{3+}] = \dots mol dm^{-3} [3]$

	(iii)	The student d	lecides to repeat t	he experiment using $0.0510\mathrm{moldm^{-3}}$ sodium thiosulfate.
		Suggest why	the student does t	his.
				[2]
(g)	Iron	is formed in st	tars by fusion reac	ctions.
	Con	nplete the nucl	ear equation for o	ne such reaction.
	48 24	+	He → '''''F	e [2]
(h)	The	table shows th	ne stable isotopes	of iron and their abundances.
	M	ass number	Abundance/	
		54	5.85	
		56	91.75	
		57	2.12	
		58	0.28	
	(i)	What is mean	t by the term mas	s number?
				[1]
	(ii)	Explain, without of iron is below		tion, how this data shows that the relative atomic mass

2	'Polybutene' is used in lip gloss.							
		e monomers for polybutene are the three isomers of butene, which are 2-methylpropene, -2-ene and isomer A .						
	(a)	Draw the skeletal formula of isomer A and name it.						
		skeletal formula:						
		name:	. [1]					
	(b)	Give the number of π bonds and σ bonds in but-2-ene, $\text{CH}_3\text{CH=CHCH}_3.$						
		π bonds σ bonds	[1]					
	(c)	But-2-ene, $CH_3CH=CHCH_3$, exists as E and Z isomers.						
		(i) Draw the structure of the E isomer of but-2-ene.						
			[1]					
		(ii) Does 2-methylpropene, $(CH_3)_2C=CH_2$, have E and Z isomers?						
		Explain your answer.						
			. [1]					
	(d)	Draw the repeating unit of the polymer of 2-methylpropene, $(CH_3)_2C=CH_2$.						
			[1]					
	(e)	The presence of the three isomers as monomers in 'polybutene' means that different side-chains are present that do not allow the polymer chains to get close together. This causes the polymer to be a liquid suitable for lip gloss.						
		Name the intermolecular bonds that hold the chains together in polybutene.						

(f) But-2-ene, $CH_3CH=CHCH_3$, reacts with hydrogen bromide.

	Dra	w the mechanism for this reaction.	
	Sho	ow curly arrows, full charges and the product.	
			[3]
(g)	But	-2-ene reacts with hydrogen to form butane.	
	Nar	me a catalyst and the corresponding conditions for this reaction.	
	cata	alyst:	
		ditions:	
			[1]
(h)	But	-2-ene is formed industrially by cracking alkanes from crude oil.	
	(i)	Complete the equation for a cracking reaction.	
		\rightarrow 2C ₄ H ₈ + C ₄ H ₁₀	[1]
	(ii)	Cracking is an example of heterogeneous catalysis.	
		Name the first step in the mechanism of heterogeneous catalysis.	
			[1]
			[.1

© OCR 2022 Turn over


3 Some students read about the reaction between hydrogen peroxide and iodide ions as shown in equation 3.1.

$$\rm 2H^+ \, + \, H_2O_2 \, + \, 2I^- \longrightarrow \, I_2 \, + \, 2H_2O \quad Equation \, 3.1$$

(a) The students find the results of an experiment which measures the concentration of hydrogen peroxide at various times.

Large excesses of acid and iodide ions are present.

The students plot a graph of these results as shown below.



(i) Use the graph to measure two half-lives to show that the kinetics are first order.

Show your working on the graph.

first half-life =

second half-life =[3]

(ii) Use the graph to measure the initial rate of the reaction in $mol dm^{-3} s^{-1}$.

initial rate = $moldm^{-3}s^{-1}$ [1]

	(iii)	The graph shows that the reaction in equation 3.1 is first order with respect to hydrogen peroxide.
		Explain why the orders with respect to the other reagents cannot be determined from this experiment.
		[2
(b)		students then set up some experiments to find the orders of reaction with respect to de and hydrogen ions.
		ne starting time they mix the hydrogen peroxide, acid and iodide ions together with a d volume of sodium thiosulfate and starch solutions.
	The form	y measure the time for the colourless solutions suddenly to go dark blue as iodine is ned.
	(i)	Suggest why the tubes go dark blue after a period of time rather than gradually from the start.
	(ii)	How can the relative rates of reaction be found from the students' results?

$$\rm 2H^+ \, + \, H_2O_2 \, + \, 2I^- \, \longrightarrow \, I_2 \, + \, 2H_2O \quad Equation \, 3.1$$

(c)* The students' results are shown below.


Use these results and information from earlier in the question to work out the rate equation for the reaction, explaining your reasoning.

Calculate the rate constant, *k*, and suggest a possible mechanism for the reaction.

Experiment	[H ₂ O ₂] /mol dm ⁻³	[I ⁻] /mol dm ⁻³	[H ⁺] /mol dm ⁻³	Initial rate of formation of I ₂ /mol dm ⁻³ s ⁻¹
1	0.030	0.010	0.10	5.25 × 10 ⁻⁶
2	0.030	0.020	0.10	1.05 × 10 ⁻⁵
3	0.030	0.020	0.20	1.05 × 10 ⁻⁵

[6]
Additional answer space if required

(d) The students find data for the variation of the rate constant, k, with temperature. From their data they plot the graph below.



Use the graph to calculate $E_{\rm a}$ in kJ ${\rm mol^{-1}}$ for the reaction in **equation 3.1**.

© OCR 2022 Turn over

		12
Chy	/motr	ypsin is an enzyme present in the small intestine.
One	e of t	ne amino acids in chymotrypsin is serine, which can be represented as HOOCCHRNH ₂ .
(a)	(i)	A molecule of serine has a chiral centre and two enantiomers.
		Draw and label two serine molecules below to illustrate the meaning of the terms chira centre and enantiomer .
		Mirror
	(ii)	[3] Glycine HOOCCH ₂ NH ₂ is another amino acid.
	(11)	
		In solution, glycine exists as a zwitterion. Draw the structure of this zwitterion.
		[1
(b)	Ser	ne reacts with glycine to form two dipeptides.
	(i)	In the boxes below, draw the structural formulae for the two dipeptides.

	(11)	explain, with a reason, whether the reaction to form the dipeptides is condensation of addition.	
(c)		motrypsin is a protein with a definite amino acid sequence and parts that have an elical structure.	
	(i)	What name is given to the amino acid sequence of a protein?	
			. [1]
	(ii)	What name is given to the $\alpha\text{-helical}$ structure and how is it held together?	

(d) Chymotrypsin catalyses the hydrolysis of proteins in the small intestine.

The active site of chymotrypsin consists of the amino acids serine, histidine and aspartic acid. These occur on different parts of the chain that are brought together by the folding of the structure.

The start of the hydrolysis mechanism is shown below. The protein being hydrolysed is at the top left.

Complete the diagram in the box below to show the results of the electron movements in step 2.

Turn over

(e) (i)	Explain why warming a solution of chymotrypsin will destroy its activity as an enzyme.						
	[2						
(ii)	Chymotrypsin has an optimum pH = 8.						
	Draw a sketch to suggest the shape of the graph of chymotrypsin activity against pH.						
	Explain the shape of your graph.						
	[3						
(iii)	A student says that the rate of an enzyme reaction is proportional to the substrate concentration.						
	Comment on the student's statement, correcting any errors.						
	[2						

	estion concerns the Advanced Notice Article 'Fluoride compounds in dental hygiene' that ed as an insert with this paper.
Нус	Profluoric acid (HF) is described in the article as 'moderately strong' with a p K_a of 3.14.
(i)	Write the expression for K_a for HF and give its units.
	K_{a} =
	units[2]
(ii)	Calculate the $\frac{[F^-]}{[HF]}$ ratio in hydrofluoric acid under the acidic conditions in plaque at pH 5.0.
	<u>[F⁻]</u> =[3]
The	e HPO ₄ ²⁻ ion is mentioned in the article.
Wri	te the formula of the conjugate base of the $HPO_4^{\ 2-}$ ion and explain your answer.
	[2]
	s said in the article to hydrolyse at 'higher pH'.
Sug	gest an equation for the reaction between SnF ₂ and hydroxide ions.
	[1]
Nar	me the shape around the N atom in the –NH ₃ ⁺ group in dectaflur.
	[1]
	(ii) The Writh

© OCR 2022 Turn over

(e)	(i)	Explain the relationship between the ion product, IP, and the solubility product, $K_{\rm sp}$, for an ionic substance.	
	(ii)	Calculate the supersaturation of calcium fluoride in saliva after brushing.	•
		Give your answer to the nearest whole number .	
		supersaturation =[3	;]
(f)	Cal	culate the ppm by mass of fluorine in pure fluoroapatite, $Ca_5(PO_4)_3F$.	
		ppm by mass =[2	<u>'</u>]

litional ar	iswer spa	ace if requ	uired				
				tional answer space if required			Tional answer space if required

END OF QUESTION PAPER

18

ADDITIONAL ANSWER SPACE

If additional space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margin(s).					



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.