



Oxford Cambridge and RSA

**GCE**

**Mathematics B MEI**

**H640/02: Pure Mathematics and Statistics**

A Level

**Mark Scheme for June 2023**

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2023

**MARKING INSTRUCTIONS****PREPARATION FOR MARKING****RM ASSESSOR**

1. Make sure that you have accessed and completed the relevant training packages for on-screen marking: *RM Assessor Online Training*; *OCR Essential Guide to Marking*.
2. Make sure that you have read and understood the mark scheme and the question paper for this unit. These are posted on the RM Cambridge Assessment Support Portal <http://www.rm.com/support/ca>
3. Log-in to RM Assessor and mark the **required number** of practice responses (“scripts”) and the **number of required** standardisation responses.

**MARKING**

1. Mark strictly to the mark scheme.
2. Marks awarded must relate directly to the marking criteria.
3. The schedule of dates is very important. It is essential that you meet the RM Assessor 50% and 100% (traditional 40% Batch 1 and 100% Batch 2) deadlines. If you experience problems, you must contact your Team Leader (Supervisor) without delay.

## 4. Annotations

Annotation	Meaning
✓ and ✗	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
E	Explanation mark 1
SC	Special case
^	Omission sign
MR	Misread
BP	Blank Page
Seen	
Highlighting	

Other abbreviations in mark scheme	Meaning
E1	Mark for explaining a result or establishing a given result
dep*	Mark dependent on a previous mark, indicated by *. The * may be omitted if only one previous M mark
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working
AG	Answer given
awrt	Anything which rounds to
BC	By Calculator
DR	This question included the instruction: In this question you must show detailed reasoning.
BP	Blank Page
Seen	
Highlighting	

## 5. Subject Specific Marking Instructions

- a. Annotations must be used during your marking. For a response awarded zero (or full) marks a single appropriate annotation (cross, tick, M0 or ^) is sufficient, but not required.

For responses that are not awarded either 0 or full marks, you must make it clear how you have arrived at the mark you have awarded and all responses must have enough annotation for a reviewer to decide if the mark awarded is correct without having to mark it independently.

It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

### Award NR (No Response)

- if there is nothing written at all in the answer space and no attempt elsewhere in the script
- OR if there is a comment which does not in any way relate to the question (e.g. 'can't do', 'don't know')
- OR if there is a mark (e.g. a dash, a question mark, a picture) which isn't an attempt at the question.

Note: Award 0 marks only for an attempt that earns no credit (including copying out the question).

If a candidate uses the answer space for one question to answer another, for example using the space for 8(b) to answer 8(a), then give benefit of doubt unless it is ambiguous for which part it is intended.

- b. An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct solutions leading to correct answers are awarded full marks but work must not always be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly. Correct but unfamiliar or unexpected methods are often signalled by a correct result following an apparently incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner.

If you are in any doubt whatsoever you should contact your Team Leader.

- c. The following types of marks are available.

### **M**

A suitable method has been selected and applied in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using

some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A method mark may usually be implied by a correct answer unless the question includes the DR statement, the command words “Determine” or “Show that”, or some other indication that the method must be given explicitly.

**A**

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

**B**

Mark for a correct result or statement independent of Method marks.

**E**

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d. When a part of a question has two or more ‘method’ steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation ‘dep\*’ is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e. The abbreviation FT implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only – differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, what is acceptable will be detailed in the mark scheme. If this is not the case please, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f. Unless units are specifically requested, there is no penalty for wrong or missing units as long as the answer is numerically correct and expressed either in SI or in the units of the question. (e.g. lengths will be assumed to be in metres unless in a particular question all the lengths are in km, when this would be assumed to be the unspecified unit.)

We are usually quite flexible about the accuracy to which the final answer is expressed; over-specification is usually only penalised where the scheme explicitly says so.

- When a value is given in the paper only accept an answer correct to at least as many significant figures as the given value.
- When a value is not given in the paper accept any answer that agrees with the correct value to 2 s.f. unless a different level of accuracy has been asked for in the question, or the mark scheme specifies an acceptable range.

NB for Specification A the rubric specifies 3 s.f. as standard, so this statement reads "3 s.f".

Follow through should be used so that only one mark in any question is lost for each distinct accuracy error.

Candidates using a value of 9.80, 9.81 or 10 for  $g$  should usually be penalised for any final accuracy marks which do not agree to the value found with 9.8 which is given in the rubric.

- g. Rules for replaced work and multiple attempts:

- If one attempt is clearly indicated as the one to mark, or only one is left uncrossed out, then mark that attempt and ignore the others.
- If more than one attempt is left not crossed out, then mark the last attempt unless it only repeats part of the first attempt or is substantially less complete.
- if a candidate crosses out all of their attempts, the assessor should attempt to mark the crossed out answer(s) as above and award marks appropriately.

- h. For a genuine misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A or B mark in the question. Marks designated as cao may be awarded as long as there are no other errors.

If a candidate corrects the misread in a later part, do not continue to follow through. E marks are lost unless, by chance, the given results are established by equivalent working. Note that a miscopy of the candidate's own working is not a misread but an accuracy error.



- i. If a calculator is used, some answers may be obtained with little or no working visible. Allow full marks for correct answers, provided that there is nothing in the wording of the question specifying that analytical methods are required such as the bold “In this question you must show detailed reasoning”, or the command words “Show” or “Determine”. Where an answer is wrong but there is some evidence of method, allow appropriate method marks. Wrong answers with no supporting method score zero. If in doubt, consult your Team Leader.
  
- j. If in any case the scheme operates with considerable unfairness consult your Team Leader.

Question		Answer	Marks	AO	Guidance
1		$\frac{a}{1-r}$ used	M1	1.1a	$a$ and $r$ are numerical values; $a = 9$ and/or $r = \pm \frac{1}{3}$
		$\frac{9}{1+\frac{1}{3}}$ <b>soi</b>	M1	1.1	
		$\frac{27}{4}$ or $6\frac{3}{4}$ or 6.75 <b>cao isw</b>	A1	1.1	if unsupported allow <b>SC2</b> for correct answer
			[3]		
2	(a)	$(x \pm 6)^2$ and $(y \pm 4)^2$	M1	1.1	completing the square twice <b>soi</b>
		$r = 7$ not from wrong working	A1	1.1	<b>NB</b> $(x - 6)^2 - 36 + (y + 4)^2 - 16 + 3 = 0$ <b>oe</b> allow <b>B2</b> for $r = 7$ unsupported
		<i>Alternatively</i> $\pm 2a = -12$ <b>oe</b> and $\pm 2b = 8$ <b>oe</b>	M1		
		$r = 7$ not from wrong working	A1		<b>NB</b> $r^2 = 6^2 + 4^2 - 3$
			[2]		
2	(b)	(6, -4)	B1	1.1	<b>FT</b> $(x \pm 6)^2 + (y \pm 4)^2$ <b>or FT</b> $\pm 2a = -12$ <b>oe</b> and $\pm 2b = 8$ <b>oe</b>
			[1]		

Question		Answer	Marks	AO	Guidance
3		take reciprocal calculate cube calculate square root to obtain $m = 343, n = 512$ <b>isw</b> or $\frac{343}{512}$ <b>isw</b>	<b>B1</b> <b>B1</b> <b>B1</b>		operations may be in any order, but 3 <b>distinct numerical</b> steps required for 3 marks if taking <b>reciprocal</b> and <b>one</b> other step are combined into one step, allow <b>B1B1B0</b>  if <b>B0B0</b> for cubing and square rooting, allow <b>SC1</b> for $\left(\sqrt{\frac{49}{64}}\right)^3 = \frac{343}{512}$ or $\left(\sqrt{\frac{64}{49}}\right)^3 = \frac{512}{343}$ seen
			[3]		
	<b>eg</b>	$\left(\frac{49}{64}\right)^{\frac{3}{2}}$ $\left(\frac{7}{8}\right)^3$ $\frac{343}{512}$ <b>isw</b>	<b>B1</b> <b>B1</b> <b>B1</b>	<b>1.1</b> <b>1.1</b> <b>1.1</b>	taking reciprocal; may be awarded after simplification square roots found; may be seen before taking reciprocal dependent on award of both preceding marks
			[3]		
	<b>eg</b>	<i>Alternatively</i> $\left(\frac{64}{49}\right)^{-3} = \left(\frac{m}{n}\right)^2$  $\left(\frac{49}{64}\right)^3 = \left(\frac{m}{n}\right)^2$  $117649 = m^2$ and $262144 = n^2$  $m = 343$ and $n = 512$	<b>B1</b> <b>B1</b> <b>B1</b>		dependent on award of both preceding marks

Question		Answer	Marks	AO	Guidance
4	(a)	$7p + 3p = 1$ oe	M1	1.2	must see some reasoning; allow explanation in words
		$p = 0.1$ or $\frac{1}{10}$ isw cao	A1	1.1	if unsupported, allow <b>SC1</b> for correct answer
			[2]		
4	(b)	$0.7$ or $\frac{7}{10}$	B1FT	1.1	their $7p$
			[1]		
4	(c)	B(30, <i>their</i> 0.3) seen or used	M1	1.1	FT their $3p$ ; allow M1 for $(\textit{their } 0.3)^2 \times (\textit{their } 0.7)^{28}$
		awrt 0.0018 isw	A1	1.1	not from wrong working; if unsupported, allow <b>SC1</b> for correct answer
			[2]		
5		$\left[ \begin{pmatrix} 5 \\ -3 \end{pmatrix} - \begin{pmatrix} 3 \\ -1 \end{pmatrix} \right] = \begin{pmatrix} 2 \\ -2 \end{pmatrix}$	B1	2.1	may be in coordinate form or may see distances identified or on diagram; may be implied by $\sqrt{(\pm 2)^2 + (\pm 2)^2}$ oe
		or $\left[ \begin{pmatrix} 3 \\ -1 \end{pmatrix} - \begin{pmatrix} 5 \\ -3 \end{pmatrix} \right] = \begin{pmatrix} -2 \\ 2 \end{pmatrix}$			
		$\sqrt{(\pm 2)^2 + (\pm 2)^2}$ oe	M1	1.1	or FT their evaluation of $\begin{pmatrix} 5 \\ -3 \end{pmatrix} - \begin{pmatrix} 3 \\ -1 \end{pmatrix}$ may be implied by correct answer
		$\sqrt{8}$ or $2\sqrt{2}$ isw	A1	1.1	if <b>B0M0</b> ; allow <b>SC1</b> for $\sqrt{80}$ or $4\sqrt{5}$ (from addition of vectors) if supported by Pythagoras;  if <b>B0M0</b> allow <b>SC1</b> for $\sqrt{8}$ or $2\sqrt{2}$ unsupported
			[3]		

Question		Answer	Marks	AO	Guidance
6		$2\cos\theta = x + 3$ or $\cos\theta = \frac{x+3}{2}$	<b>B1</b>	<b>2.1</b>	allow sign errors in their expressions for $\sin\theta$ and $\cos\theta$ ; allow if just see brackets expanded, but must be 3 terms in each case
		$2\sin\theta = y - 1$ or $\sin\theta = \frac{y-1}{2}$	<b>B1</b>	<b>1.1</b>	
		$\left(\frac{x\pm 3}{2}\right)^2 + \left(\frac{y\pm 1}{2}\right)^2 = \cos^2\theta + \sin^2\theta$ or $(x \pm 3)^2 + (y \pm 1)^2 = 4\cos^2\theta + 4\sin^2\theta$ oe	<b>M1</b>	<b>1.1</b>	
		$(x + 3)^2 + (y - 1)^2 = 4$	<b>A1</b>	<b>1.1</b>	
			<b>[4]</b>		
		<i>Alternatively</i> centre of circle is $(-3, 1)$	<b>B1</b>		allow one sign error in bracket;
		radius is 2	<b>B1</b>		
		$(x + 3)^2 + (y - 1)^2 = 2^2$	<b>M1</b>		
		$(x + 3)^2 + (y - 1)^2 = 4$	<b>A1</b>		
7		$(2x)^8$ <b>soi</b>	<b>M1</b>	<b>3.1a</b>	allow recovery from bracket error; may be implied by award of second <b>M1</b>
		${}^{12}C_8$ or ${}^{12}C_4$ or 495 seen	<b>B1</b>	<b>1.1</b>	allow <b>M1</b> for $495 \times 2 \times k^4 = 79\,200\,000$ <b>oe</b> ; “ = 79 200 000” may be implied by $k = 5$
		$495 \times 256 \times k^4 [x^8] = 79\,200\,000 [x^8]$ <b>oe</b>	<b>M1</b>	<b>1.1</b>	
		$k = 5$ <b>cao isw</b>	<b>A1</b>	<b>3.2a</b>	not from wrong working; but allow recovery from $x^8$ on one side of equation only
			<b>[4]</b>		allow <b>SC2</b> for $k = 5$ unsupported
			<b>[4]</b>		

Question		Answer	Marks	AO	Guidance
8	(a)	<b>not</b> a simple random sample since <b>each possible sample</b> does <b>not</b> have an <b>equal probability</b> of being selected	<b>B1</b>	<b>2.4</b>	allow “ <b>not</b> a simple random sample since <b>each plant</b> does <b>not</b> have an <b>equal probability</b> of being selected” ignore further comments unless contradictory  the essential elements of the comment are in bold
			[1]		
8	(b)	18 (shorter than 40 cm) seen 110 (shorter than 80 cm) <b>or</b> 10 (taller than 80 cm) seen $\frac{92}{120}$ <b>oe or</b> $\frac{28}{120}$ <b>oe</b>  <b>awrt</b> 77% > 75% <b>oe</b> so this supports the (owner’s) statement	<b>B1</b> <b>B1</b> <b>M1</b>  <b>A1</b>	<b>2.1</b> <b>1.1</b> <b>1.1</b>  <b>2.2a</b>	may be embedded in calculation may be embedded in calculation <b>FT</b> their 18 and their 110; allow <b>M1</b> for $\frac{91}{120}$ <b>or</b> $\frac{29}{120}$ <b>oe</b>  <b>or</b> $\frac{75}{100} \times 120 = 90$ <b>oe</b> <b>or</b> 92 > 90 so this supports the (owner’s) statement; need comparison <b>and</b> comment
			[4]		
8	(c)	must refer to sample in answer  eg since different <b>samples</b> give different results, not all samples would necessarily support owner’s statement  eg no since another <b>sample</b> may contain all the tallest plants (and/or shortest)  eg no since the heights in other <b>samples</b> may be different	<b>B1</b>	<b>2.4</b>	ignore superfluous comments referring to eg growing conditions, soil type etc  do not allow eg no, conditions different for other plants  eg no, different plants on different rows  eg no, might be biggest and smallest plants on other rows  eg the first sample is not representative
			[1]		

Question		Answer	Marks	AO	Guidance
9	(a)	eg sample, since only some data used <b>oe</b> eg sample, since data for 2009 not used <b>oe</b>	<b>B1</b>	<b>2.2a</b>	sample, since not all data used; ignore further reasoning unless contradictory
			[1]		
9	(b)	£27 085 or £27 100 or £27 000	<b>B1</b>	<b>3.4</b>	
			[1]		
9	(c)	it's interpolation <b>oe</b>  eg because we use 2009 <b>oe</b> which is between 2008 and 2010  eg a straight line model appears to be justified  eg relationship seems to be linear  eg median income seems to be directly proportional to time  eg positive correlation [between median income and year]	<b>B1</b>       <b>B1</b>	<b>2.2a</b>      <b>2.2b</b>	do not allow  eg 27085 lies between 2008 and 2010  eg 27085 lies between the values for 2008 and 2010  do not allow  eg positive association  eg positive relationship
			[2]		


Question		Answer	Marks	AO	Guidance
9	(d)	<p><b>not reasonable oe</b> since eg (median taxable) income varies considerably across different regions of London eg (median) income different in Croydon and Camden eg (median) income lower in Croydon (than Camden) eg pattern of change in income over time different in Camden to Croydon</p>	<b>B1</b>	<b>2.4</b>	<b>LDS advantage</b>
			[1]		
10		<p><math>4x \times \frac{1}{2} \sin 2x - \int \frac{1}{2} \sin 2x \times 4 dx</math> <math>2x \sin 2x - \int 2 \sin 2x dx</math> <b>oe</b> <math>F[x] = 2x \sin 2x + \cos 2x</math> <math>F\left[\frac{\pi}{4}\right] - F[0]</math> <math>\frac{\pi}{2} - 1</math> <b>or</b> <math>\frac{\pi-2}{2}</math> <b>isw</b></p>	<p><b>M1*</b> <b>A1</b> <b>A1</b> <b>M1dep*</b> <b>A1</b></p>	<p><b>3.1a</b> <b>1.1</b> <b>1.1</b> <b>1.1</b> <b>3.2a</b></p>	<p>allow sign errors only, condone omission of dx allow omission of dx; may be unsimplified ignore + c must see substitution if F[x] incorrect, otherwise may be implied by correct answer; allow recovery from bracket error</p>
			[5]		
11	(a)	<p><math>\frac{dy}{dx} = k\sqrt{x}</math> <math>3 = k \times \sqrt{4}</math> <math>k = \frac{3}{2}</math> <b>or</b> <math>\frac{dy}{dx} = \frac{3}{2}\sqrt{x}</math> <b>isw</b></p>	<p><b>B1</b> <b>M1</b> <b>A1</b></p>	<p><b>2.1</b> <b>1.1</b> <b>2.2a</b></p>	<p>may be implied by final answer if <b>B0M0</b> allow <b>SC1</b> for <math>\frac{dy}{dx} = \frac{k}{\sqrt{x}}</math> and <math>k = 6</math> as final answer <b>or</b> <math>\frac{dy}{dx} = \frac{6}{\sqrt{x}}</math> as final answer</p>
			[3]		



Question		Answer	Marks	AO	Guidance
11	(b)	<p>their <math>k \times \frac{x^{\frac{3}{2}}}{\frac{3}{2}}</math> <b>oe</b></p> <p><math>10 = (\sqrt{4})^3 + c</math></p> <p><math>c = 2</math> or <math>y = x^{\frac{3}{2}} + 2</math> or <math>y = x\sqrt{x} + 2</math> isw</p>	<p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p>	<p><b>3.1a</b></p> <p><b>1.1</b></p> <p><b>1.1</b></p>	<p><b>FT</b> their <math>k</math></p> <p><b>FT</b> their integration, one term in <math>x</math> with index 1.5</p> <p>must see 'y =' at some point</p> <p>if <b>B0M0</b> allow <b>SC2</b> for <math>y = 12x^{\frac{1}{2}} - 14</math> or <math>y = 12\sqrt{x} - 14</math></p> <p>or <math>y = 12x^{\frac{1}{2}} + c</math> and <math>c = -14</math></p>
			[3]		
12	(a)	<p>because it's neither a one-to-one nor a many-to-one (mapping)</p> <p>do not allow eg because it's neither a one-to-one nor a many-to-one <b>function</b></p>	<b>B1</b>	<b>2.4</b>	<p>allow because it's one-to-many (mapping)</p> <p>allow eg because each value of <math>x</math> is mapped to two values <b>oe</b></p> <p>do not allow eg because it's a one-to-many <b>function</b></p>
			[1]		
12	(b)	<p><math>\frac{x^2+2}{x^2+2-3}</math></p> <p><math>\frac{x^2+2}{x^2-1}</math> or <math>\frac{x^2+2}{(x-1)(x+1)}</math></p>	<p><b>M1</b></p> <p><b>A1</b></p>	<p><b>1.1</b></p> <p><b>1.1</b></p>	
			[2]		

Question		Answer	Marks	AO	Guidance
12	(c)	$ x  > 1$ <b>or</b> $x < -1$ <b>or</b> $x > 1$ <b>or</b> $x < -1, x > 1$ <b>or</b> $x < -1 \cup x > 1$	<b>B1</b>	<b>1.1</b>	do not allow eg $x < -1$ and $x > 1$ eg $-1 > x > 1$
			[1]		
13	(a)	$H_0: \mu = 0.14$ $H_1: \mu < 0.14$  their $\mu$ is the <b>population mean mass</b> of this variety of apple	<b>B1</b>  <b>B1</b>	<b>1.1</b>  <b>2.5</b>	allow any other symbol except $\bar{x}$ or $\bar{X}$ , as long as it is correctly defined; allow hypotheses stated in words  allow weight;  correct definition of $\mu$ may be embedded in hypotheses written out as a sentence;  do not allow $\bar{x}$ or $\bar{X}$
			[2]		
13	(b)	$[\bar{X} \sim] N\left(0.14, \frac{0.0199^2}{80}\right)$	<b>B1</b>  <b>B1</b>	<b>3.3</b>  <b>2.2a</b>	Normal distribution with correct mean or variance allow variance = <b>awrt</b> $4.95 \times 10^{-6}$ or <b>awrt</b> $0.00222^2$  all correct, but allow full credit if no symbol used; allow symbol other than $\bar{X}$ if correctly defined as sample mean, but do not allow $\mu$
			[2]		
13	(c)	<b>awrt</b> 0.136 seen <b>BC</b>	<b>B1</b>	<b>1.1</b>	
		$\bar{X} < 0.136$ only or $\bar{X} \leq 0.136$ only	<b>B1</b>	<b>3.4</b>	<b>FT</b> other correctly defined symbol
			[2]		

Question		Answer	Marks	AO	Guidance
13	(d)	0.1316 < 0.136 <b>or</b> 0.1316 is in the critical region (must be correct critical region) <b>oe</b>	<b>M1</b>	<b>3.4</b>	condone $p = \text{awrt } 0.00007 < 0.05$ <b>oe NB</b> 0.0000740 <b>or</b> $z = \text{awrt } -3.79 < -1.645$ <b>oe</b> from use of $\bar{X} \sim N\left(0.14, \frac{0.0198^2}{80}\right)$
		<b>or</b> $p = \text{awrt } 0.00008 < 0.05$ <b>oe</b>			
		<b>NB</b> 0.0000799			
		<b>or</b> $z = \text{awrt } -3.78 < -1.645$ <b>oe</b>			
		reject $H_0$	<b>A1</b>	<b>1.1</b>	allow accept $H_1$ <b>or</b> result is significant
		there is sufficient evidence at the 5% level to <b>suggest</b> that the <b>mean</b> mass of the apples is <b>less than</b> 0.14 kg	<b>A1</b>	<b>2.2b</b>	allow weight; do not allow eg conclude / prove / indicate or other assertive statement instead of suggest
			[3]		
14	(a)	discard City of London (as part of the data not available)	<b>B1</b>	<b>2.4</b>	<b>LDS advantage</b> do not allow if answer spoiled
		<b>or</b> discard any regions where one or more pieces of data are missing <b>oe</b>			eg because it's an anomaly, eg because it's an outlier,
			[1]		
14	(b)	<b>scatter</b> does not look linear <b>oe</b>	<b>B1</b>	<b>3.4</b>	ignore extra comments unless they contradict an otherwise correct answer
		pmcc not close to 1 <b>oe</b>	<b>B1</b>	<b>3.4</b>	ignore extra comments unless they contradict an otherwise correct answer
			[2]		

Question		Answer	Marks	AO	Guidance
14	(c)	$27216 \pm 2 \times 4177.5$ or $61.0 \pm 2 \times 5.32$  $m < 18861$ or $m > 35\,571$  percentage $< 50.36$ or percentage $> 71.64$  <p style="text-align: center;">Scatter diagram to show Percentage of Pupils Achieving 5 A*-C Grades against Median Income of Taxpayers</p> 	<b>M1</b>  <b>A1</b>  <b>A1</b>  <b>A1</b>	<b>1.1</b>  <b>1.1</b>  <b>1.1</b>  <b>1.1</b>	use of 2 standard deviation check for one of the 4 calculations <b>soi</b>  allow $\leq$ and $\geq$  allow $\leq$ and $\geq$  if <b>M1A0A0</b> allow <b>M1 SCB1</b> for all 4 correct values seen
			[4]		
14	(d)	between 0 and 0.3743 since  eg outliers gave a false impression of linearity  eg scatter will be more like a circle	<b>B1</b>	<b>2.4</b>	need to refer to the shape of the scatter <b>oe</b>
			[1]		

Question		Answer	Marks	AO	Guidance
15		$y = 1$ then $x = 2$ only  $\frac{1}{y} \times \frac{dy}{dx}$ $x^3 \times \frac{dy}{dx} + 3x^2y$  $\frac{1}{y} \times \frac{dy}{dx} + x^3 \times \frac{dy}{dx} + 3x^2y [= 0]$  substitution of their $x = 2$ <b>and</b> $y = 1$ to obtain numerical value for $\frac{dy}{dx}$  $y - 1 = \left(\text{their } \frac{3}{4}\right)(x - \text{their } 2)$ <b>oe</b>  $3x - 4y - 2 = 0$ or $-3x + 4y + 2 = 0$ <b>oe</b>	<b>B1</b>  <b>B1</b> <b>M1</b>  <b>A1</b>  <b>M1*</b>  <b>M1dep*</b>  <b>A1</b>	<b>3.1a</b>  <b>2.1</b> <b>1.1</b>  <b>1.1</b>  <b>1.1</b>  <b>1.1</b>  <b>3.1a</b>  <b>3.2a</b>	first term correct; allow $y'$ for $\frac{dy}{dx}$  Product Rule; allow one coefficient error <b>or</b> one index error      <b>NB</b> $-\frac{4}{3}$  dependent on at least two of 3 terms correct on LHS following differentiation; if expression for $\frac{dy}{dx}$ or evaluation of $\frac{dy}{dx}$ is incorrect, need to see substitution for award of <b>M1</b>    <b>FT</b> negative reciprocal of their $-\frac{4}{3}$ and their 2 may see eg $1 = \frac{3}{4} \times 2 + c$  must be in required form, but coefficients may be fractions
			[7]		

Question		Answer	Marks	AO	Guidance
16	(a)	0.16×0.84×2	M1	1.1	condone omission of 2
		or B(2, 0.16) or B(2, 0.84) seen or $1 - (0.84^2 + 0.16^2)$ $\frac{168}{625}$ or 0.2688 or 0.269 or 0.27 <b>cao</b>	A1	1.1	allow recovery from bracket error mark the final answer allow <b>SC1</b> for correct answer unsupported
			[2]		
16	(b)	$0.75 - 0.66 = 0.09$	M1	3.1a	allow 0.09 embedded in correct place in Venn diagram or contingency table; allow <b>M1</b> for 9%
		[0.16 – 0.09 = ] 0.07 <b>isw</b>	A1	1.1	allow <b>SC1</b> for correct answer unsupported
			[2]		
16	(c)	$\frac{0.09}{0.75}$	M1	3.1a	<b>M0</b> for 0.12 from wrong working
		0.12	A1	1.1	allow <b>SC1</b> for correct answer unsupported
			[2]		
16	(d)	$0.12 \neq 0.16$	M1	2.1	
		so not independent	A1	2.2a	if <b>M0</b> allow <b>SCB1</b> for $0.16 \times 0.75 \neq 0.09$ so <b>not</b> independent
			[2]		

Question		Answer	Marks	AO	Guidance
17		divide through by $\cos x$ to obtain	<b>B1</b>	<b>2.1</b>	
		$2\tan x + \sec^2 x = 4$			
		$2\tan x + \tan^2 x + 1 = 4$	<b>M1*</b>	<b>3.1a</b>	use of Pythagoras to obtain equation in $\tan x$ only; allow 1 sign error
		$\tan^2 x + 2\tan x - 3 [= 0]$	<b>A1</b>	<b>1.1</b>	
		$\tan x = 1$ or $-3$	<b>M1*dep</b>	<b>1.1</b>	2 values obtained for $\tan x$ from their quadratic
		$[x =] -1.24905$ to $-1.249$ or $-1.25$ or $-1.2$			
		$[x =] 1.8925$ to $1.893$ or $1.89$ or $1.9$	<b>A1</b>	<b>3.2a</b>	any two correct
	$[x =] \frac{\pi}{4}$ or $0.785$ to $0.7854$ or $0.79$				
	$[x =] -\frac{3\pi}{4}$ or $-2.3562$ to $-2.356$ or $-2.36$ or $-2.4$	<b>A1</b>	<b>2.2a</b>	all four correct and <b>no extra</b> values in range; ignore <b>correct</b> extra values outside range but <b>A0</b> if incorrect values outside range	
			<b>[6]</b>		

Question	Answer	Marks	AO	Guidance
	<p><i>alternatively</i> multiply through by <math>\cos x</math> to obtain</p> $2\sin x \cos x + 1 = 4\cos^2 x$ $\sin 2x + 1 = 2\cos 2x + 2$ $5\cos^2 2x + 4\cos 2x [= 0]$ <p><b>NB</b> square both sides: <math>\sin^2 2x = 4\cos^2 2x + 4\cos 2x + 1</math> <b>oe</b></p> $\cos 2x = 0 \text{ or } -0.8$ <p>2 values obtained for <math>\cos 2x</math> from their quadratic</p> <p>[x =] -1.24905 to -1.249 or -1.25 or -1.2</p> <p>[x =] 1.8925 to 1.893 or 1.89 or 1.9</p> <p>[x =] <math>\frac{\pi}{4}</math> or 0.785 to 0.7854 or 0.79</p> <p>[x =] <math>-\frac{3\pi}{4}</math> or -2.3562 to -2.356 or -2.36 or -2.4</p>	<p><b>B1</b></p> <p><b>M1*</b></p> <p><b>A1</b></p> <p><b>M1dep*</b></p> <p><b>A1</b></p> <p><b>A1</b></p>		<p>use of double angle formulae, allow 1 sign error</p> <p>or <math>\sqrt{5}\cos(2x + 0.4636 \dots) = -1</math></p> <p>or <math>\sqrt{5}\sin(2x - 1.1071 \dots) = 1</math></p> <p><math>\cos(2x + 0.4636 \dots) = -\frac{1}{\sqrt{5}}</math> <b>or</b> <math>\sin(2x - 1.1071 \dots) = \frac{1}{\sqrt{5}}</math></p> <p>any two correct</p> <p>all four correct and <b>no extra</b> values in range; ignore <b>correct</b> extra values outside range but <b>A0</b> if incorrect values outside range</p>



Question		Answer	Marks	AO	Guidance
18	(a)	260 cao	B1	1.1	
			[1]		
18	(b)	31 cao	B1	1.1	mark the final answer
			[1]		
18	(c)	any 2 distinct reasons eg (approximately) symmetrical (about the mean)  eg approximately bell-shaped / unimodal  eg data is continuous	B1  B1	3.5a  3.5a	ignore extra comments unless they contradict an otherwise correct answer
			[2]		
18	(d)	[variance is] awrt 62.2  or [sd is] 7.89 seen BC    0.26287 – 0.263134 or 0.26	M1    A1	3.3    3.4	NB 62.15567...may be implied by sd = 7.88 or 7.89  NB 0.263047...from $\sqrt{62.2}$ or 0.263133... from 7.89  NB 0.262871... from 7.88, 0.262973...from unrounded sd  allow B2 for correct answer unsupported
			[2]		
18	(e)	B(28, $p$ ) used, where $p$ is value calculated in (d)  $0.888 \leq p < 0.896$	M1  A1	3.1a  1.1	may be given to 2 sf; allow B2 for correct answer unsupported
			[2]		

Question		Answer	Marks	AO	Guidance
18	(f)	$7.8 + 0.18 \times 260$	M1	3.1a	
		$0.18^2 \times 62.2$ oe	M1	3.5c	or $0.18 \times \sqrt{62.2}$
		$N(54.6, 2.0138 - 2.02)$	A1	1.1	allow eg $1.42^2$ for variance
			[3]		

APPENDIX

Exemplar responses for Q2(b)

Response	Mark

## Need to get in touch?

If you ever have any questions about OCR qualifications or services (including administration, logistics and teaching) please feel free to get in touch with our customer support centre.

### Call us on

**01223 553998**

### Alternatively, you can email us on

**support@ocr.org.uk**

### For more information visit



**ocr.org.uk/qualifications/resource-finder**



**ocr.org.uk**



**Twitter/ocrexams**



**/ocrexams**



**/company/ocr**



**/ocrexams**



**CAMBRIDGE**  
UNIVERSITY PRESS & ASSESSMENT

OCR is part of Cambridge University Press & Assessment, a department of the University of Cambridge.

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored. © OCR 2023 Oxford Cambridge and RSA Examinations is a Company Limited by Guarantee. Registered in England. Registered office The Triangle Building, Shaftesbury Road, Cambridge, CB2 8EA.

Registered company number 3484466. OCR is an exempt charity.

OCR operates academic and vocational qualifications regulated by Ofqual, Qualifications Wales and CCEA as listed in their qualifications registers including A Levels, GCSEs, Cambridge Technicals and Cambridge Nationals.

OCR provides resources to help you deliver our qualifications. These resources do not represent any particular teaching method we expect you to use. We update our resources regularly and aim to make sure content is accurate but please check the OCR website so that you have the most up-to-date version. OCR cannot be held responsible for any errors or omissions in these resources.

Though we make every effort to check our resources, there may be contradictions between published support and the specification, so it is important that you always use information in the latest specification. We indicate any specification changes within the document itself, change the version number and provide a summary of the changes. If you do notice a discrepancy between the specification and a resource, please [contact us](#).

Whether you already offer OCR qualifications, are new to OCR or are thinking about switching, you can request more information using our [Expression of Interest form](#).

Please [get in touch](#) if you want to discuss the accessibility of resources we offer to support you in delivering our qualifications.